arXiv:1002.4704v2 [math.AT] 2 Mar 2010

REAL BOTT MANIFOLDS AND ACYCLIC DIGRAPHS

SUYOUNG CHOI AND SANG-IL OUM

Abstract. Masuda (2008) provided the characterization of real Bott manifolds in terms of three operations on upper triangular matrices. We provide a combinatorial characterization of real Bott manifolds up to diffeomorphism in terms of operations on directed acyclic graphs. Our observation leads to several new invariants of real Bott manifolds.

1. Introduction

A *small cover*, defined by Davis and Januszkiewicz [DJ91], is an n-dimensional closed smooth manifold M with a smooth action of real torus $(S^0)^n (=: T)$ satisfying the following two conditions:

- the action is locally isomorphic to a standard action of T on $\mathbb{R}^n$, and
- the orbit space M/T can be identified with a simple (combinatorial) polytope P.

A *cube* is a polytope combinatorially equivalent to the cartesian product of finitely many intervals. We will restrict our attention to the case where P is a cube. In this case, a small cover is said to be *over* a cube.

Small covers over cubes are known as *real Bott manifolds*, see [CMS10] or Section 3. They are obtained as iterated $\mathbb{R}P^1$ bundles starting with a point, where each stage is the projectivization of a Whitney sum of two real line bundles. The topological classification of real Bott manifolds is known by Kamishima and Masuda [KM09], and Masuda [Mas08]; two real Bott manifolds are diffeomorphic if and only if their cohomology rings with $\mathbb{Z}_2$ coefficients are isomorphic as graded rings. Choi [Cho08b] showed that small covers over cubes are strongly related to acyclic digraphs.

We introduce two operations on acyclic digraphs and show that all diffeomorphism types of real Bott manifolds can be described by the composition of these two operations. This combinatorial observation allows us to invent several invariants of real Bott manifolds.

Our combinatorial interpretation allows more efficient enumeration of all n-dimensional real Bott manifolds up to diffeomorphism. We list the number B_n of n-dimensional real Bott manifolds in Table 1 for $n \leq 8$. Previously, B_n were computed only for $n \leq 5$ in [Mas08] and [Naz08], but it was a hard task even for

---

Date: March 2, 2010.

2000 *Mathematics Subject Classification.* Primary 37F20, 57R91, 05C90; Secondary 53C25.

Key words and phrases. small cover over cube, real Bott manifold, acyclic digraph, invariant, local complementation, Bott equivalence.

The first author was supported by the Japanese Society for the Promotion of Sciences (JSPS grant no. P09023). The second author was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2009-0063183) and TJ Park Junior Faculty Fellowship.

n	1	2	3	4	5	6	7	8
B_n	1	2	4	12	54	472	8,512	328,416
E_n	1	1	2	3	8	29	222	3,607
S_n		1		2		6		31

TABLE 1. The numbers B_n, E_n, S_n of n-dimensional real Bott manifolds, orientable real Bott manifolds and symplectic real Bott manifolds up to diffeomorphism, respectively.

$n = 5$ when using topological or linear algebraic methods. The classification for 8-dimensional real Bott manifolds takes less than 10 minutes on a regular desktop computer, when using the list of non-isomorphic acyclic digraphs provided by B. D. McKay.[1] In addition, we also list the numbers E_n and S_n of n-dimensional orientable real Bott manifolds and symplectic real Bott manifolds in Table 1 for $n \leq 8$, respectively.

The rest of this paper is organized as follows. In Sections 2 and 3, we briefly review some notions in acyclic digraphs and small covers respectively. In Section 4, we classify the acyclic digraph up to Bott equivalence, which implies the diffeomorphism class of real Bott manifolds. Finally, in Section 5, we introduce several invariants of acyclic digraphs up to Bott equivalence.

2. ACYCLIC DIGRAPHS

2.1. **Preliminaries on acyclic digraphs.** We briefly review the terminologies in graph theory. A *directed graph*, or simply a *digraph* D is a pair (V, E) consisting of a finite set V of *vertices* and a set E of ordered pairs $e = (u, v)$ of distinct vertices of D, called *arcs*. We write $V(D)$ to denote the set of all vertices of D. An ordering $v_1, v_2, \ldots, v_n$ of vertices of a digraph D is an *acyclic ordering* if $i < j$ for each arc (v_i, v_j) in D. A digraph is *acyclic* if it admits an acyclic ordering.

A vertex u is *adjacent* to a vertex v in D if (u, v) is an arc of D. If (u, v) is an arc of D, then v is called an *out-neighbor* of u and u is called an *in-neighbor* of v. We write $N_D^+(v)$, $N_D^-(v)$ to denote the set of all out-neighbors and in-neighbors of v, respectively, in D. The *in-degree* of a vertex v in D, denoted by $\deg_D^-(v)$, is the number of in-neighbors of v. Similarly the *out-degree* of a vertex v in D, denoted by $\deg_D^+(v)$, is the number of out-neighbors of v.

For two digraphs $D = (V, E)$ and $D' = (V', E')$, a bijection $f : V \to V'$ is called an *isomorphism* when $(u, v) \in E$ if and only if $(f(u), f(v)) \in E'$. Two digraphs are *isomorphic* if there is an isomorphism.

For a digraph $D = (V, E)$ with a fixed ordering $\{v_1, v_2, \ldots, v_n\}$ of V, the *adjacency matrix* of D is an $n \times n$ matrix $A_D = (a_{ij})_{i,j \in \{1,2,\ldots,n\}}$ such that

$$a_{ij} = \begin{cases} 1 & \text{if } (v_i, v_j) \in E, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\mathcal{G}_n$ be the set of all acyclic digraphs on the vertex set $\{1, 2, \ldots, n\}$ and let $\mathcal{A}_n$ be the set of all adjacency matrices of acyclic digraphs in $\mathcal{G}_n$.

---

[1] `http://cs.anu.edu.au/~bdm/data/digraphs.html`



FIGURE 1. Two operations: a local complementation and a slide

2.2. **Bott equivalence of acyclic digraphs.** An acyclic digraph H is *Bott equivalent* to an acyclic digraph D if H has an isomorphic digraph that is obtained from D by successively applying local complementations and slides. In the following, we define local complementations and slides.

For a vertex v of a digraph D, let $D * v$ be a digraph obtained by adding an arc (u, w) if $(u, w) \notin E$, or removing the arc (u, w) otherwise, for each pair $(u, w) \in N_D^-(v) \times N_D^+(v)$ with $u \neq w$. This operation to obtain $D * v$ from D is called a *local complementation* at v. This operation is not new; As far as the authors find, Bouchet [Bou87] wrote one of the earliest papers on local complementations on digraphs.

For two sets X and Y, we write $X \Delta Y = (X \setminus Y) \cup (Y \setminus X)$. For two distinct vertices v, w having the same set of in-neighbors in a digraph D, we define $D\Delta vw$ to be a digraph obtained by replacing $N_D^+(w)$ by $N_D^+(w)\Delta N_D^+(v)$. This operation to obtain $D\Delta vw$ from D is called a *slide*. See Figure 1.

It is easy to observe that if D is an acyclic digraph, then so are $D * v$ and $D\Delta xy$ (assuming $N_D^-(x) = N_D^-(y)$). Furthermore $D * v * v = D$ and $D\Delta xy\Delta xy = D$.

3. SMALL COVERS OVER CUBES

3.1. **Small covers over cubes.** Let P be a simple polytope of dimension n and let $\mathcal{F}(P) = \{F_1, \ldots, F_m\}$ be the set of facets, codimension one faces, of P. It is known by [DJ91] that all small covers over P can be classified by certain maps $\lambda : \mathcal{F}(P) \to \mathbb{Z}_2^n$ satisfying the so-called *non-singularity condition*; $\{\lambda(F_{i_1}), \ldots, \lambda(F_{i_n})\}$ is a basis of $\mathbb{Z}_2^n$ whenever the intersection $F_{i_1} \cap \cdots \cap F_{i_n}$ is non-empty. We call λ a *characteristic function*.

The construction of the small cover with respect to λ is simple. Let $S^0(F_i)$ be the subgroup of $T(= (S^0)^n)$ generated by $\lambda(F_i)$. Given a point $p \in P$, we denote by $G(p)$ the minimal face containing p in its relative interior. Assume $G(p) = F_{j_1} \cap \cdots \cap F_{j_k}$. Then $S^0(G(p)) = \oplus_{i=1}^k S^0(F_{j_i})$. Note that $S^0(G(p))$ is a k-dimensional subgroup of T. Let $M(\lambda)$ denote

$$M(\lambda) = P \times T / \sim, \tag{1}$$

where $(p, g) \sim (q, h)$ if $p = q$ and $g^{-1}h \in S^0(G(p))$. The canonical free action of T on $P \times T$ descends to an action on $M(\lambda)$. It is easy to show that the action is locally standard and the orbit space can be identified with P. Thus $M(\lambda)$ is a small cover over P.

Two small covers M and N over P are *Davis-Januskiewicz equivalent* (or simply, *D-J equivalent*) if there is a weak T-equivariant homeomorphism $f : M \to N$ covering the identity on P. By [DJ91], $M(\lambda_1)$ is D-J equivalent to $M(\lambda_2)$ if and only if there is an automorphism $\sigma \in \mathrm{Aut}(\mathbb{Z}_2^n)$ such that $\lambda_1 = \sigma \circ \lambda_2$.

When P is an n-dimensional cube, P has $2n$ facets, and for each facet F of P there exists a unique facet which does not intersect with F. So we give an ordering of the facets such that the facets F_j and F_{n+j} do not intersect for $1 \leq j \leq n$. Then we may assign an $n \times 2n$ matrix to a characteristic function λ by choosing a basis for $(\mathbb{Z}_2)^n$. Then

$$(\lambda(F_1) \cdots \lambda(F_{2n})) = (A|B),$$

where A and B are $n \times n$ matrices. Since $\bigcap_{j=1}^{n} F_j \neq \emptyset$, we also may assume that A is the identity matrix of size n. Then we can see that the non-singularity condition of λ is equivalent to the following; *every principal minor of B is 1.*

Hence, there is a bijection between the DJ-equivalence classes of small covers over cubes and the set $M(n)$ of $n \times n$ matrices over $\mathbb{Z}_2$ all of whose principal minors are 1.

Theorem 3.1 (Choi [Cho08b, Theorem 2.2])**.** *Let $\phi : \mathcal{A}_n \to M(n)$ by $A \mapsto I_n + A^t$, where I_n is the identity matrix of size n and A^t is the transpose matrix of A. Then ϕ is a bijection.*

Let $D \in \mathcal{G}_n$ and let A_D be its adjacency matrix. By the above theorem, $(I_n|\phi(A_D))$ is the characteristic function. For the sake of simplicity, we denote the corresponding small cover by $M(A_D)$.

Let $D, H \in \mathcal{G}_n$. We note that the ordering of vertices of an acyclic digraph is nothing but the ordering of facets of P, and the diffeomorphism type of the small cover is independent on the ordering of facets by the construction (1). Hence, if D and H are isomorphic, then $M(A_D)$ and $M(A_H)$ are diffeomorphic (but not D-J equivalent).

3.2. **Real Bott manifolds.** To define a real Bott manifold, we consider with a sequence of $\mathbb{R}P^1$ bundles

$$M_n \xrightarrow{\pi_n} M_{n-1} \xrightarrow{\pi_{n-1}} \cdots \xrightarrow{\pi_2} M_1 \xrightarrow{\pi_1} M_0 = \{\text{a point}\}, \tag{2}$$

where each fiber bundle $\pi_j : M_j \to M_{j-1}$ for $j = 1, \ldots, n$ is the projectivization of a Whitney sum of a real line bundle and the trivial real line bundle over M_{j-1}. We call a sequence (2) a *real Bott tower* and M_n a *real Bott manifold*. It is well-known that real line bundles are classified by their first Stiefel-Whitney classes and $H^1(M_{j-1}; \mathbb{Z}_2)$ is isomorphic to $(\mathbb{Z}_2)^{j-1}$. Hence, each M_j is determined by a vector A_j in $(\mathbb{Z}_2)^{j-1}$. One can organize this vector into an $n \times n$ upper-triangular square matrix A with zero diagonals by putting A_j as the upper part of the j-th column vector.

Proposition 3.2 (Panov and Masuda [MP08, Proposition 3.1])**.** *Let A be an upper-triangular square matrix with zero diagonals. A real Bott tower with respect to A carries a natural real torus action turning it into a small cover $M(A)$*[2].

If D is an acyclic digraph, then its adjacency matrix A_D with respect to its acyclic ordering is an upper-triangular square matrix with zero diagonals. Hence, all small covers over a cube are diffeomorphic to real Bott manifolds.

---

[2] The original statement of [MP08, Proposition 3.1] is written in terms of complex versions of a Bott tower and a small cover called a *quasitoric manifold*. But the same argument can be easily applied to the real cases.

4. Classification of real Bott manifolds

For a matrix A, we write A_j to denote the j-th column of A and write A_j^i to denote its (i,j)-th entry of A. Masuda [Mas08] defined three operations Φ_S, Φ^k and Φ_C^I on $\mathcal{A}_n$ as follows: (Notice that these matrices are over $\mathbb{Z}_2$ and therefore $1+1=0$.)

(1) $\Phi_S(A) = SAS^{-1}$, where S is a permutation matrix

(2) For $k \in \{1,\ldots,n\}$,

$$\Phi^k(A)_j = A_j + A_j^k A_k \quad \text{for } j = 1,\ldots,n.$$

(3) Let $I = \{i_1, i_2, \ldots, i_\ell\}$ be a subset of $\{1,\ldots,n\}$ such that $A_i = A_j$ for $i,j \in I$ and $i_1 < i_2 < \cdots < i_\ell$. Let C be an invertible $|I| \times |I|$ matrix over $\mathbb{Z}_2$. Then,

$$\Phi_C^I(A)_j^i = \begin{cases} \sum_{k=1}^{|I|} C_k^m A_j^{i_k} & \text{if } i = i_m \in I, \\ A_j^i & \text{if } i \notin I. \end{cases}$$

Roughly speaking, Φ_C^I is the left matrix multiplication of C to the submatrix of A which consists of i-th rows for $i \in I$.

Remark 4.1. Indeed, Masuda [Mas08] described all three operations on square upper-triangular matrices over $\mathbb{Z}_2$. But since every small cover over a cube is diffeomorphic to a real Bott manifold, these three operations can be extended to $\mathcal{A}_n$. Moreover, this is more natural because the upper-triangular matrices are not closed under Φ_S.

We say that two matrices in $\mathcal{A}_n$ are *Bott equivalent* if one is transformed to the other through a sequence of the three operations. Masuda [Mas08] proved that Bott equivalence on $\mathcal{A}_n$ characterizes real Bott manifolds up to diffeomorphisms.

Theorem 4.2 (Masuda [Mas08, Theorem 1.1]). *Two matrices $A, B \in \mathcal{A}_n$ are Bott equivalent if and only if corresponding real Bott manifolds $M(A)$ and $M(B)$ are diffeomorphic.*

We now prove our main observation, stating that two acyclic digraphs D and H are Bott equivalent (in the sense of the definition in Section 2) if and only if A_D and A_H are Bott equivalent.

Lemma 4.3. *Two acyclic digraphs are Bott equivalent if and only if their adjacency matrices are Bott equivalent.*

Proof. Let $D = (V,E)$ be an acyclic digraph with a certain ordering $v_1, v_2, \ldots, v_n$ of the vertices. Obviously $A_{D * v_k} = \Phi^k(A_D)$ and if $i < j$, then $A_{D \Delta v_i v_j} = \Phi_C^I(A_D)$ with $C = \left(\begin{smallmatrix} 1 & 0 \\ 1 & 1 \end{smallmatrix}\right)$. If $i > j$, then $A_{D \Delta v_i v_j} = \Phi_C^I(A_D)$ with $C = \left(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}\right)$. Therefore Bott equivalent acyclic digraphs have Bott equivalent adjacency matrices.

Now, let us prove the converse. We want to show that each of three operations Φ_S, Φ^k, Φ_C^I on the adjacency matrix A_D of D gives an adjacency matrix of another acyclic digraph Bott equivalent to D. Firstly, Φ_S will simply give an isomorphic digraph, which is again Bott equivalent to D. Secondly, $\Phi^k(A_D) = A_{D * v_k}$ and therefore Φ^k corresponds to a local complementation at v_k.

Last, we claim that Φ_C^I can be expressed as a sequence of slides. Suppose that this claim is false. Since C is invertible, there is a sequence of elementary row operations $\sigma_1, \sigma_2, \ldots, \sigma_t$ so that $C = \sigma_1 \circ \sigma_2 \circ \cdots \circ \sigma_t(I)$. Since C is a matrix over

$\mathbb{Z}_2$, each σ_i is either an operation to interchange two rows of C or an operation to add a row to another row in C. Let $C' = \sigma_2 \circ \cdots \circ \sigma_t(I)$. Then C' is invertible and $C = \sigma_1(C')$. We assume that C is chosen as a counterexample with minimum t. Thus, $\Phi^I_{C'}(A_D) = A_{D'}$ for some acyclic digraph D' Bott equivalent to D. If σ_1 is an operation to interchange two rows, then $\Phi^I_{C'}(A_{D'})$ is an adjacency matrix after swapping two corresponding vertices in the fixed order, because these two vertices have the same set of in-neighbors and therefore clearly $\Phi^I_{C'}(A_{D'})$ is an adjacency matrix of a digraph isomorphic to D'.

If σ_1 is an operation to add a row to another, then we can simulate σ_1 by a slide on D' and therefore $\Phi^I_C(A_D) = A_{D'\Delta uv}$ for two vertices $u, v \in I$. This proves the claim because $D'\Delta uv$ is Bott equivalent to D. □

Now we are ready to see the following theorem, implied by Theorem 4.2 and Lemma 4.3.

Theorem 4.4. *Two acyclic digraphs D and H are Bott equivalent if and only if the small covers over cubes $M(A_D)$ and $M(A_H)$ are diffeomorphic.*

5. Invariants of real Bott manifolds

In this section, we introduce topological invariants of small covers over cubes. To do so, we focus on finding parameters on acyclic digraphs that are invariant under taking local complementations and slides.

A vertex in an acyclic digraph is called a *root* if it has no in-neighbor. Let $L_k(D)$ be the set of vertices v such that a longest directed path ending at v has exactly k edges. Each $L_k(D)$ is called the k-th level of D. Clearly $L_0(D)$ is the set of roots. For an acyclic digraph D with n vertices, a *level sequence* of D is a sequence $(|L_0(D)|, |L_1(D)|, |L_2(D)|, \ldots, |L_{n-1}(D)|)$. It is obvious that $\cup_{i=0}^{n-1} L_i(D) = V(D)$. Note that for each vertex $v \in L_i(D)$, all in-neighbors of v are in $\cup_{j=0}^{i-1} L_j(D)$ and all out-neighbors of v are in $\cup_{j=i+1}^{n-1} L_j(D)$.

Proposition 5.1. *Bott equivalent acyclic digraphs have the identical level sequence.*

Proof. Let D be an acyclic digraph. It is enough to show that both local complementations and slides do not change $L_i(D)$. Let v be a vertex in $L_i(D)$. Then there is a longest directed path P_v in D ending at v with exactly i edges.

Let us first show that $v \in L_i(D * w)$ for a vertex w in D. It is enough to show that P_v is a path in $D * w$ as well, because, if so, then a longest path in $D * w$ ending at v will be a path in $D * w * w = D$ as well. Suppose that P_v is not a path in $D * w$. Then w must remove at least one arc (x, y) of P_v and therefore (x, w) and (w, y) are arcs of D. Since D is acyclic, w is not on P_v. Then by replacing the arc (x, y) by a path xwy in P_v, we can find a path longer than P_v in D, contradictory to the assumption that P_v is a longest path ending at v. This proves the claim that $L_i(D) = L_i(D * w)$ for all i.

Now let us prove that $v \in L_i(D\Delta uw)$ for two vertices u, w having the set of in-neighbors. Again, it is enough to show that $D\Delta uw$ has a path of length i ending at v; because if $D\Delta uw$ has a longer path ending at v, then so does D by the fact that $(D\Delta uw)\Delta uw = D$. We may assume that $D\Delta uw$ removes at least one arc (x, y) of P_v. Then $w = x$ and both (w, y) and (u, y) are arcs of D. Since u and w have the same set of in-neighbors, we can replace w by u in P_v to obtain a path of the same length in $D\Delta uw$. This completes the proof. □

The *rank* of a digraph D is the rank of the adjacency matrix A_D over $\mathbb{Z}_2$.

Proposition 5.2. *Bott equivalent acyclic digraphs have the same rank.*

Proof. Let D be an acyclic digraph. For a vertex v, we can obtain A_{D*v} from A_D by adding the row of v to rows of in-neighbors of v. Note that since D is acyclic, no in-neighbor of v is an out-neighbor of v. Therefore $\operatorname{rank}(A_{D*v}) = \operatorname{rank}(A_D)$.

Similarly let u, w be two distinct vertices of D having the same set of in-neighbors. Clearly D and $D\Delta uw$ have the same rank because $A_{D\Delta uw}$ is obtained by adding the row of u to the row of w in A_D and therefore $\operatorname{rank}(A_{D\Delta uw}) = \operatorname{rank}(A_D)$. □

For subsets X and Y of the vertex set of a digraph D, we write $A_D[X, Y]$ to denote the submatrix of A_D whose rows correspond to X and columns correspond to Y. We write $\rho_D(X, Y) = \operatorname{rank}(A_D[X, Y])$ where $A_D[X, Y]$ is considered as a matrix over $\mathbb{Z}_2$. We write $\rho_D(X) = \rho_D(X, V(D) \setminus X)$.

Proposition 5.3. *Let D, H be Bott equivalent acyclic digraphs on n vertices. Then,*

(i) $\rho_D(\cup_{i\in I} L_i(D)) = \rho_H(\cup_{i\in I} L_i(H))$ *for every subset* I *of* $\{0, 1, 2, \ldots, n-1\}$,
(ii) $\rho_D(L_i(D), L_{i+1}(D)) = \rho_H(L_i(H), L_{i+1}(H))$ *for all* $i \in \{0, 1, 2, \ldots, n-2\}$.

Proof. It is enough to prove when $H = D * v$ or $H = D\Delta uw$. By Proposition 5.1, $L_i(D) = L_i(H)$ for each i.

(i) For a subset $I = \{0, 1, 2, \ldots, n-1\}$, let $X = \cup_{i\in I} L_i(D)$ and $Y = V(D) \setminus X$. Let $M = A_D[X, Y]$ and $M' = A_H[X, Y]$. Then $\rho_D(X) = \operatorname{rank} M$ and $\rho_H(X) = \operatorname{rank} M'$.

Let us first consider the case when $H = D * v$ for a vertex v of D. Then either $v \in X$ or $v \in Y$. If $v \in X$, then M has a row indexed by v and M' is obtained from M by adding the row of v to all rows indexed by in-neighbors of v in X. If $v \in Y$, then M has a column indexed by v and M' is obtained from M by adding the column of v to all columns indexed by out-neighbors of v in Y. Thus, in both cases, $\operatorname{rank} M = \operatorname{rank} M'$.

Now let us consider the case when $H = D\Delta uw$ for two vertices u, w having the same set of in-neighbors. Since u and w have the same set of in-neighbors, they belong to the same level. Therefore either $\{u, w\} \subseteq X$ or $\{u, w\} \subseteq Y$. If $\{u, w\} \subseteq Y$, then $M' = M$. If $\{u, w\} \subseteq X$, then M' is obtained from M by adding the row of u to the row of w. So $\operatorname{rank} M = \operatorname{rank} M'$. This completes the proof of (i).

(ii) Since D is acyclic, there is no arc from $L_a(D)$ to $L_b(D)$ if $a > b$. Therefore

$$\rho_D(L_i(D), L_{i+1}(D)) = \rho_D(L_i(D) \cup L_{i+2}(D) \cup L_{i+3}(D) \cup \cdots \cup L_{n-1}(D)),$$

and by (i), we have $\rho_D(L_i(D), L_{i+1}(D)) = \rho_H(L_i(H), L_{i+1}(H))$. □

For two vertices x and y of D, we say that $x \sim_D y$ if x and y have the same set of in-neighbors. Then $\sim_D$ is an equivalence relation of $V(D)$ and each equivalence class is called a *sibling group* of D. A sibling group of D corresponds to a maximal set of identical columns in A_D. Clearly each level $L_i(D)$ is partitioned into sibling groups.

Proposition 5.4. *If D and H are Bott equivalent acyclic digraphs, then $L_i(D)$ and $L_i(H)$ have the same number of sibling groups having exactly k vertices for each pair of i and k.*

Proof. We may assume that either $H = D * v$ for a vertex v or $H = D\Delta uw$ for two vertices u, w having the same set of in-neighbors.

We claim that $\sim_D$ and $\sim_H$ are identical equivalence relations. To do so, we will prove that if x and y have the same set of in-neighbors in D, then they have the same set of in-neighbors in H.

Firstly let us consider the case when $H = D * v$ for a vertex v. We may assume that (v, x) or (v, y) is an arc of D, because otherwise the set of in-neighbors of x or y will not be changed by applying a local complementation at v. Since x and y have the same set of in-neighbors, both (v, x) and (v, y) are arcs of D. Then x and y still have the same set of in-neighbors in $D * v$.

Secondly let us assume that $H = D\Delta uw$. Similarly we may assume that both (u, x) and (u, y) are arcs of D. Since x and y have the same set of in-neighbors, (w, x) is an arc of D if and only if (w, y) is an arc of D. After a slide, (w, x) is an arc of H if and only if (w, x) is not an arc of D and similarly (w, y) is an arc of H if and only if (w, y) is not an arc of D. This proves the claim. □

The *odd height* of an acyclic digraph D is the length of a longest directed path ending at a vertex of odd out-degree. In other words, it is the maximum k such that L_k contains a vertex of odd out-degree. If D has no vertex of odd out-degree, then we assume that its odd height is ∞.

Proposition 5.5. *Bott equivalent acyclic digraphs have the same odd height.*

Proof. Let D be an acyclic digraph. Let v be a vertex of D. Let x, y be two distinct vertices having the same set of in-neighbors. Then

$$\deg^+_{D*v}(w) \equiv \begin{cases} \deg^+_D(w) + \deg^+_D(v) & \pmod 2 \quad \text{if } w \text{ is an in-neighbor of } v, \\ \deg^+_D(w) & \pmod 2 \quad \text{otherwise.} \end{cases}$$

$$\deg^+_{D\Delta xy}(w) \equiv \begin{cases} \deg^+_D(w) + \deg^+_D(x) & \pmod 2 \quad \text{if } w = y, \\ \deg^+_D(w) & \pmod 2 \quad \text{otherwise.} \end{cases}$$

If every vertex of D has even out-degree, then clearly a local complementation and a slide do not create a vertex of odd out-degree. So we may assume that D has a vertex of odd out-degree. Let k be the odd height of D. Let w be a vertex of odd out-degree in $L_k(D)$.

Let us first consider $D * v$ for a vertex $v \in L_i(D)$. If w is an in-neighbor of v in D, then the out-degree of w in $D * v$ is odd because $i > k$ and $\deg^+_D(v)$ is even. If w is not an in-neighbor of v in D, then the out-degree of w in $D * v$ is again odd. This proves that the odd height of $D * v$ is at least the odd height of D. Since $(D * v) * v = D$, we conclude that D and $D * v$ have the same odd height.

Now we claim that $L_k(D\Delta xy)$ has a vertex of odd out-degree in $D\Delta xy$. Suppose that the out-degree of w in $D\Delta xy$ is even. Then $w = y$, $x \in L_k(D)$, and the out-degree of x in D is odd. Since $(D\Delta xy)\Delta xy = D$, we conclude that D and $D\Delta xy$ have the same odd height. □

We observe that the above invariants have topological motivations. For an acyclic digraph D, let $M(A_D)$ be the small cover corresponding to D. Choi [Cho08a] showed that $M(A_D)$ is orientable if and only if the out-degree of every vertex of D is even; in other words, the odd height of D is ∞. Hence, the odd height can be seen as a generalization of the orientability of real Bott manifolds. In addition, Ishida [Ish10] proved that $M(A_D)$ is symplectic if and only if $V(D)$ can be partitioned



FIGURE 2. Theses acyclic digraphs are not Bott equivalent but have the identical set of invariants discussed in this paper

into pairs of vertices so that two vertices in a pair have the same set of in-neighbors. Equivalently, the cardinality of each sibling group is even. Hence the number of sibling groups with a fixed size is also a generalization of the symplecticness of real Bott manifolds. It is easy to see that if the cardinality of each sibling group is even, then its odd height must be ∞. It is obvious with topological viewpoint as well, because every symplectic manifold is orientable. We computed a few values of the numbers of n-dimensional orientable real Bott manifolds and symplectic manifolds up to diffeomorphism in Table 1.

We remark that the invariants discussed in this paper completely classify all Bott equivalence classes up to 4 vertices but not on 5 vertices. One can easily check that two acyclic digraphs in Figure 2 are not Bott equivalent but have the same set of invariants.

ACKNOWLEDGMENT

We would like to thank B. D. McKay. His computer program called *nauty* to test graph isomorphism has been very useful for our enumeration. We also used the list of non-isomorphic acyclic digraphs up to 8 vertices posted on his website.

REFERENCES

[Bou87] André Bouchet, *Digraph decompositions and Eulerian systems*, SIAM J. Algebraic Discrete Methods **8** (1987), no. 3, 323–337. MR 897733 (88k:05087)

[Cho08a] Suyoung Choi, *The number of orientable small covers over cubes*, arXiv:0812.3861 (2008).

[Cho08b] ———, *The number of small covers over cubes*, Algebr. Geom. Topol. **8** (2008), no. 4, 2391–2399. MR 2465745 (2009i:57072)

[CMS10] Suyoung Choi, Mikiya Masuda, and Dong Youp Suh, *Quasitoric manifolds over a product of simplices*, Osaka J. Math. **47** (2010), no. 1, 1–21.

[DJ91] Michael W. Davis and Tadeusz Januszkiewicz, *Convex polytopes, Coxeter orbifolds and torus actions*, Duke Math. J. **62** (1991), no. 2, 417–451. MR 1104531 (92i:52012)

[Ish10] Hiroaki Ishida, *Symplectic real Bott manifolds*, arXiv:1001.3726 (2010).

[KM09] Yoshinobu Kamishima and Mikiya Masuda, *Cohomological rigidity of real Bott manifolds*, Algebr. Geom. Topol. **9** (2009), 2479–2502.

[MP08] Mikiya Masuda and Taras E. Panov, *Semi-free circle actions, Bott towers, and quasitoric manifolds*, Mat. Sb. **199** (2008), no. 8, 95–122. MR 2452268 (2009i:57073)

[Mas08] Mikiya Masuda, *Classification of real Bott manifolds*, arXiv:0809.2178 (2008).

[Naz08] Admi Nazra, *Real Bott tower*, Tokyo Metropolitan University, Master Thesis (2008).

Department of Mathematics, Osaka City University, Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan
E-mail address: `choi@sci.osaka-cu.ac.jp`

Department of Mathematical Sciences, KAIST, 335 Gwahangno, Yuseong-gu, Daejeon 305-701, Republic of Korea
E-mail address: `sangil@kaist.edu`